

HOMEWORK 4

Due date: Feb 6, 2020, 11:55PM.

Bibliography: Trench Chap. 4. The first exercise in each problem set is solved for you to use as a model.

1. Exercises 1-5, p. 138

Ex 1. Half life $\tau = 3200$ years, initial amount $Q_0 = 20$ g. Radioactive decay model $Q' = -kQ$, $k = (\ln 2)/\tau$ has solution

$$Q(t) = e^{-kt}Q_0 = \exp\left[-\frac{t}{\tau} \ln 2\right] Q_0 = 20 \times 2^{-t/\tau} \text{ g.}$$

Ex 2.

Ex 3.

Ex 4.

Ex 5.

2. Exercises 15-19, pp. 138-9

Ex 15. Gold creation rate $r = 1$ oz/hr, theft rate $s = \frac{1}{20}W(t)$ oz/hr. Model equation $W' = r - s = r - \frac{1}{20}W$, and solution to homogeneous problem $W' + \frac{1}{20}W = 0$ is $W_h = e^{-t/20}$. Variation of parameters, $W = uW_h$

$$u'W_h = r \Rightarrow u' = re^{t/20} \Rightarrow u = 20re^{t/20} + c \Rightarrow W(t) = 20r + ce^{-t/20}.$$

Initial condition $W(0) = 20r + c = 1 \Rightarrow c = 1 - 20r \Rightarrow W(t) = 20r(1 - e^{-t/20}) + e^{-t/20}$, $\lim_{t \rightarrow \infty} W(t) = 20r$.

Check in Maxima

```
(%i3) ode2('diff(W,t)=r-W/20,W,t);
```

```
(%o3) W = e^(-t/20) (20 r e^(t/20) + %c)
```

```
(%i4) ic1(%,t=0,W=1);
```

```
(%o4) W = e^(-t/20) (20 r e^(t/20) - 20 r + 1)
```

```
(%i5)
```

Ex 16.

Ex 17.

Ex 18.

Ex 19.

3. Exercises 1-5, p.148

Ex 1. Room temperature $T_0 = 70^\circ \text{F}$, freezer temperature $T_1 = 12^\circ \text{F}$. Model equation for thermometer temperature $T(t)$ is $T' = -k(T - T_1)$, $T(0) = T_0$. Solution is $T = T_1 + (T_0 - T_1)e^{-kt}$. From $T(1/2) = 40$ obtain

$$12 + 58e^{-k/2} = 40 \Rightarrow k = -2 \ln \frac{28}{58} = \ln \left(\frac{29}{14} \right)^2.$$

Evaluate

$$T(2) = 12 + 58 \exp \left[-2 \ln \left(\frac{29}{14} \right)^2 \right] = 12 + 58 \left(\frac{14}{29} \right)^4 = 15.15^\circ \text{F}.$$

```
(%i13) float(12+58*(14/29)^4)
```

```
(%o13) 15.15027266390586
```

(%i14)

Ex 2.

Ex 3.

Ex 4.

Ex 5.

4. Exercises 1-5, p. 160

Ex 1. Weight $m = 192 \text{ lb} = 87.3 \text{ kg}$, model equation is $mv' = mg - kv$, with $g = 9.8 \text{ m/s}^2$, $k = 2.5 \text{ lbf s/ft} = 36.54 \text{ kg/s}$, with solution $v(t) = mg/k [\exp(-kt/m) - 1]$, and terminal velocity $\lim_{t \rightarrow \infty} v(t) = mg/k = 23.4 \text{ m/s}$.

Ex 2.

Ex 3.

Ex 4.

Ex 5.