

- Recall: to each matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$ associate four fundamental subspaces:
 - Column space**, $C(\mathbf{A}) = \{\mathbf{b} \in \mathbb{R}^m \mid \exists \mathbf{x} \in \mathbb{R}^n \text{ such that } \mathbf{b} = \mathbf{A}\mathbf{x}\} \subseteq \mathbb{R}^m$, the part of \mathbb{R}^m reachable by linear combination of columns of \mathbf{A}
 - Left null space**, $N(\mathbf{A}^T) = \{\mathbf{y} \in \mathbb{R}^m \mid \mathbf{A}^T \mathbf{y} = \mathbf{0}\} \subseteq \mathbb{R}^m$, the part of \mathbb{R}^m not reachable by linear combination of columns of \mathbf{A}
 - Row space**, $R(\mathbf{A}) = C(\mathbf{A}^T) = \{\mathbf{c} \in \mathbb{R}^n \mid \exists \mathbf{y} \in \mathbb{R}^m \text{ such that } \mathbf{c} = \mathbf{A}^T \mathbf{y}\} \subseteq \mathbb{R}^n$, the part of \mathbb{R}^n reachable by linear combination of rows of \mathbf{A}
 - Null space**, $N(\mathbf{A}) = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{A}\mathbf{x} = \mathbf{0}\} \subseteq \mathbb{R}^n$, the part of \mathbb{R}^n not reachable by linear combination of rows of \mathbf{A}

The fundamental theorem of linear algebra (FTLA) states

$$C(\mathbf{A}), N(\mathbf{A}^T) \leq \mathbb{R}^m, C(\mathbf{A}) \perp N(\mathbf{A}^T), C(\mathbf{A}) \cap N(\mathbf{A}^T) = \{\mathbf{0}\}, C(\mathbf{A}) \oplus N(\mathbf{A}^T) = \mathbb{R}^m$$

$$C(\mathbf{A}^T), N(\mathbf{A}) \leq \mathbb{R}^n, C(\mathbf{A}^T) \perp N(\mathbf{A}), C(\mathbf{A}^T) \cap N(\mathbf{A}) = \{\mathbf{0}\}, C(\mathbf{A}^T) \oplus N(\mathbf{A}) = \mathbb{R}^n$$

- New concepts:
 - Proof techniques, properties of direct sums

Lemma 1. Let \mathcal{U}, \mathcal{V} , be subspaces of vector space \mathcal{W} . Then $\mathcal{W} = \mathcal{U} \oplus \mathcal{V}$ if and only if

i. $\mathcal{W} = \mathcal{U} + \mathcal{V}$, and

ii. $\mathcal{U} \cap \mathcal{V} = \{\mathbf{0}\}$.


Proof. $\mathcal{W} = \mathcal{U} \oplus \mathcal{V} \Rightarrow \mathcal{W} = \mathcal{U} + \mathcal{V}$ by definition of direct sum, sum of vector subspaces. To prove that $\mathcal{W} = \mathcal{U} \oplus \mathcal{V} \Rightarrow \mathcal{U} \cap \mathcal{V} = \{\mathbf{0}\}$, consider $w \in \mathcal{U} \cap \mathcal{V}$. Since $w \in \mathcal{U}$ and $w \in \mathcal{V}$ write

$$w = w + \mathbf{0} \quad (w \in \mathcal{U}, \mathbf{0} \in \mathcal{V}), \quad w = \mathbf{0} + w \quad (\mathbf{0} \in \mathcal{U}, w \in \mathcal{V}),$$

and since expression $w = u + v$ is unique, it results that $w = \mathbf{0}$. Now assume (i),(ii) and establish an unique decomposition. Assume there might be two decompositions of $w \in \mathcal{W}$, $w = u_1 + v_1$, $w = u_2 + v_2$, with $u_1, u_2 \in \mathcal{U}$, $v_1, v_2 \in \mathcal{V}$. Obtain $u_1 + v_1 = u_2 + v_2$, or $x = u_1 - u_2 = v_2 - v_1$. Since $x \in \mathcal{U}$ and $x \in \mathcal{V}$ it results that $x = \mathbf{0}$, and $u_1 = u_2$, $v_1 = v_2$, i.e., the decomposition is unique. \square

Lemma 2. Orthogonal complements of \mathbb{R}^m ($m \in \mathbb{N}$, finite), $\mathcal{U}, \mathcal{V} \leq \mathbb{R}^m$, $\mathcal{U} = \mathcal{V}^\perp$, $\mathcal{V} = \mathcal{U}^\perp$, form a direct sum $\mathcal{U} \oplus \mathcal{V} = \mathbb{R}^m$. (proved after discussion of Gram-Schmidt procedure)

FTLA - Graphical representation



$$\mathbb{R}^n = C(A^T) \oplus N(A)$$

$$C(A^T) \perp N(A)$$

usually: $m \geq n$

$$\mathbb{R}^m = N(A^T) \oplus C(A)$$

$$N(A^T) \perp C(A)$$

i. $C(\mathbf{A}) \leq \mathbb{R}^m$ (column space is vector subspace of codomain of $\mathbf{A}: \mathbb{R}^n \rightarrow \mathbb{R}^m$)

Proof. Consider arbitrary $\mathbf{u}, \mathbf{v} \in C(\mathbf{A})$, $\alpha, \beta \in \mathbb{R}$. Verify vector subspace properties (Lesson 7 p.4):

i. Inclusion (elements of $C(\mathbf{A})$ are in \mathbb{R}^m) $\mathbf{u} \in \mathbb{R}^m$, yes by definition of $C(\mathbf{A}) = \{\mathbf{b} \in \mathbb{R}^m \mid \exists \mathbf{x} \in \mathbb{R}^n \text{ such that } \mathbf{b} = \mathbf{A}\mathbf{x}\}$. (This immediately results from definitions and will not be shown explicitly in following proofs).

ii. Closed ($\alpha\mathbf{u} + \beta\mathbf{v} \in C(\mathbf{A})$). By definition of $C(\mathbf{A})$, $\mathbf{u}, \mathbf{v} \in C(\mathbf{A})$ implies existence of $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ such that $\mathbf{u} = \mathbf{A}\mathbf{x}$, $\mathbf{v} = \mathbf{A}\mathbf{y}$. Compute $\alpha\mathbf{u} + \beta\mathbf{v} = \alpha(\mathbf{A}\mathbf{x}) + \beta(\mathbf{A}\mathbf{y}) = \mathbf{A}(\alpha\mathbf{x} + \beta\mathbf{y})$, and note that since $\alpha\mathbf{x} + \beta\mathbf{y} \in \mathbb{R}^n$, $\alpha\mathbf{u} + \beta\mathbf{v} \in C(\mathbf{A})$. \square

ii. $N(\mathbf{A}^T) \leq \mathbb{R}^n$ (left null space is vector subspace of domain of $\mathbf{A}: \mathbb{R}^n \rightarrow \mathbb{R}^m$)

Proof. Consider arbitrary $\alpha, \beta \in \mathbb{R}$, $\mathbf{x}, \mathbf{y} \in N(\mathbf{A}^T) \Rightarrow \mathbf{A}^T\mathbf{x} = \mathbf{0}$, $\mathbf{A}^T\mathbf{y} = \mathbf{0}$. Compute $\mathbf{A}^T(\alpha\mathbf{x} + \beta\mathbf{y}) = \alpha(\mathbf{A}^T\mathbf{x}) + \beta(\mathbf{A}^T\mathbf{y}) = \alpha \cdot \mathbf{0} + \beta \cdot \mathbf{0} = \mathbf{0}$, hence $\alpha\mathbf{x} + \beta\mathbf{y} \in N(\mathbf{A}^T)$ \square

iii. $C(\mathbf{A}) \perp N(\mathbf{A}^T)$ (column space is orthogonal to left null space).

Proof. Consider arbitrary $\mathbf{u} \in C(\mathbf{A})$, $\mathbf{v} \in N(\mathbf{A}^T)$. By definition of $C(\mathbf{A})$, $\exists \mathbf{x} \in \mathbb{R}^n$ such that $\mathbf{u} = \mathbf{A}\mathbf{x}$, and by definition of $N(\mathbf{A}^T)$, $\mathbf{A}^T\mathbf{v} = \mathbf{0}$. Compute $\mathbf{u}^T\mathbf{v} = (\mathbf{A}\mathbf{x})^T\mathbf{v} = \mathbf{x}^T\mathbf{A}^T\mathbf{v} = \mathbf{x}^T(\mathbf{A}^T\mathbf{v}) = \mathbf{x}^T\mathbf{0} = \mathbf{0}$, hence $\mathbf{u} \perp \mathbf{v}$ for arbitrary \mathbf{u}, \mathbf{v} , and $C(\mathbf{A}) \perp N(\mathbf{A}^T)$. \square

iv. $C(\mathbf{A}) \cap N(\mathbf{A}^T) = \{\mathbf{0}\}$ ($\mathbf{0}$ is the only vector both in $C(\mathbf{A})$ and $N(\mathbf{A}^T)$).

Proof. (By contradiction, *reductio ad absurdum*). Assume there might be $\mathbf{b} \in C(\mathbf{A})$ and $\mathbf{b} \in N(\mathbf{A}^T)$ and $\mathbf{b} \neq \mathbf{0}$. Since $\mathbf{b} \in C(\mathbf{A})$, $\exists \mathbf{x} \in \mathbb{R}^n$ such that $\mathbf{b} = \mathbf{A}\mathbf{x}$. Since $\mathbf{b} \in N(\mathbf{A}^T)$, $\mathbf{A}^T\mathbf{b} = \mathbf{A}^T(\mathbf{A}\mathbf{x}) = \mathbf{0}$. Note that $\mathbf{x} \neq \mathbf{0}$ since $\mathbf{x} = \mathbf{0} \Rightarrow \mathbf{b} = \mathbf{0}$, contradicting assumptions. Multiply equality $\mathbf{A}^T\mathbf{A}\mathbf{x} = \mathbf{0}$ on left by \mathbf{x}^T ,

$$\mathbf{x}^T\mathbf{A}^T\mathbf{A}\mathbf{x} = \mathbf{0} \Rightarrow (\mathbf{A}\mathbf{x})^T(\mathbf{A}\mathbf{x}) = \mathbf{b}^T\mathbf{b} = \|\mathbf{b}\|^2 = \mathbf{0},$$

thereby obtaining $\mathbf{b} = \mathbf{0}$, using norm property 3 (Lesson 4, p5). Contradiction.

\square

$$\text{v. } C(\mathbf{A}) \oplus N(\mathbf{A}^T) = \mathbb{R}^m$$

Proof. (iii) and (iv) have established that $C(\mathbf{A}), N(\mathbf{A}^T)$ are orthogonal complements

$$C(\mathbf{A}) = N(\mathbf{A}^T)^\perp, N(\mathbf{A}^T) = C(\mathbf{A})^\perp.$$

By Lemma 2 it results that $C(\mathbf{A}) \oplus N(\mathbf{A}^T) = \mathbb{R}^m$. (Reminder: Proof of Lemma 2 is postponed until discussion of the Gram-Schmidt procedure). \square

The remainder of the FTLA is established by considering $\mathbf{B} = \mathbf{A}^T$, e.g., since it has been established in (v) that $C(\mathbf{B}) \oplus N(\mathbf{A}^T) = \mathbb{R}^n$, replacing $\mathbf{B} = \mathbf{A}^T$ yields $C(\mathbf{A}^T) \oplus N(\mathbf{A}) = \mathbb{R}^m$, etc.

Remark. The great widespread applicability of linear algebra results in large part due to the complete characterization of the possible solutions to $\mathbf{A}\mathbf{x} = \mathbf{b}$ provided by the FTLA and the orthogonal decomposition of the domain and codomain.