

DATA INFORMATION

1. Partition of linear mapping domain and codomain

A partition of a set S has been introduced as a collection of subsets $P = \{S_i | S_i \subset P, S_i \neq \emptyset\}$ such that any given element $x \in S$ belongs to only one set in the partition. This is modified when applied to subspaces of a vector space, and a partition of a set of vectors is understood as a collection of subsets such that any vector except $\mathbf{0}$ belongs to only one member of the partition.

Linear mappings between vector spaces $f: U \rightarrow V$ can be represented by matrices A with columns that are images of the columns of a basis $\{\mathbf{u}_1, \mathbf{u}_2, \dots\}$ of U

$$A = [f(\mathbf{u}_1) \ f(\mathbf{u}_2) \ \dots].$$

Consider the case of real finite-dimensional domain and co-domain, $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$, in which case $A \in \mathbb{R}^{m \times n}$,

$$A = [f(\mathbf{e}_1) \ f(\mathbf{e}_2) \ \dots \ f(\mathbf{e}_n)] = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n].$$

The column space of A is a vector subspace of the codomain, $C(A) \leq \mathbb{R}^m$, but according to the definition of dimension if $n < m$ there remain non-zero vectors within the codomain that are outside the range of A ,

$$n < m \Rightarrow \exists \mathbf{v} \in \mathbb{R}^m, \mathbf{v} \neq \mathbf{0}, \mathbf{v} \notin C(A).$$

All of the non-zero vectors in $N(A^T)$, namely the set of vectors orthogonal to all columns in A fall into this category. The above considerations can be stated as

$$C(A) \leq \mathbb{R}^m, \ N(A^T) \leq \mathbb{R}^m, \ C(A) \perp N(A^T) \ C(A) + N(A^T) \leq \mathbb{R}^m.$$

The question that arises is whether there remain any non-zero vectors in the codomain that are not part of $C(A)$ or $N(A^T)$. The fundamental theorem of linear algebra states that there no such vectors, that $C(A)$ is the orthogonal complement of $N(A^T)$, and their direct sum covers the entire codomain $C(A) \oplus N(A^T) = \mathbb{R}^m$.

LEMMA 1. Let \mathcal{U}, \mathcal{V} , be subspaces of vector space \mathcal{W} . Then $\mathcal{W} = \mathcal{U} \oplus \mathcal{V}$ if and only if

- i. $\mathcal{W} = \mathcal{U} + \mathcal{V}$, and
- ii. $\mathcal{U} \cap \mathcal{V} = \{\mathbf{0}\}$.

Proof. $\mathcal{W} = \mathcal{U} \oplus \mathcal{V} \Rightarrow \mathcal{W} = \mathcal{U} + \mathcal{V}$ by definition of direct sum, sum of vector subspaces. To prove that $\mathcal{W} = \mathcal{U} \oplus \mathcal{V} \Rightarrow \mathcal{U} \cap \mathcal{V} = \{\mathbf{0}\}$, consider $\mathbf{w} \in \mathcal{U} \cap \mathcal{V}$. Since $\mathbf{w} \in \mathcal{U}$ and $\mathbf{w} \in \mathcal{V}$ write

$$\mathbf{w} = \mathbf{w} + \mathbf{0} \quad (\mathbf{w} \in \mathcal{U}, \mathbf{0} \in \mathcal{V}), \quad \mathbf{w} = \mathbf{0} + \mathbf{w} \quad (\mathbf{0} \in \mathcal{U}, \mathbf{w} \in \mathcal{V}),$$

and since expression $\mathbf{w} = \mathbf{u} + \mathbf{v}$ is unique, it results that $\mathbf{w} = \mathbf{0}$. Now assume (i),(ii) and establish an unique decomposition. Assume there might be two decompositions of $\mathbf{w} \in \mathcal{W}$, $\mathbf{w} = \mathbf{u}_1 + \mathbf{v}_1$, $\mathbf{w} = \mathbf{u}_2 + \mathbf{v}_2$, with $\mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}$, $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{V}$. Obtain $\mathbf{u}_1 + \mathbf{v}_1 = \mathbf{u}_2 + \mathbf{v}_2$, or $\mathbf{x} = \mathbf{u}_1 - \mathbf{u}_2 = \mathbf{v}_2 - \mathbf{v}_1$. Since $\mathbf{x} \in \mathcal{U}$ and $\mathbf{x} \in \mathcal{V}$ it results that $\mathbf{x} = \mathbf{0}$, and $\mathbf{u}_1 = \mathbf{u}_2$, $\mathbf{v}_1 = \mathbf{v}_2$, i.e., the decomposition is unique. \square

In the vector space $U + V$ the subspaces U, V are said to be orthogonal complements is $U \perp V$, and $U \cap V = \{\mathbf{0}\}$. When $U \leq \mathbb{R}^m$, the orthogonal complement of U is denoted as U^\perp , $U \oplus U^\perp = \mathbb{R}^m$.

THEOREM. Given the linear mapping associated with matrix $A \in \mathbb{R}^{m \times n}$ we have:

1. $C(A) \oplus N(A^T) = \mathbb{R}^m$, the direct sum of the column space and left null space is the codomain of the mapping
2. $C(A^T) \oplus N(A) = \mathbb{R}^n$, the direct sum of the row space and null space is the domain of the mapping
3. $C(A) \perp N(A^T)$ and $C(A) \cap N(A^T) = \{\mathbf{0}\}$, the column space is orthogonal to the left null space, and they are orthogonal complements of one another,

$$C(A) = N(A^T)^\perp, \ N(A^T) = C(A)^\perp.$$

4. $C(A^T) \perp N(A)$ and $C(A^T) \cap N(A) = \{\mathbf{0}\}$, the row space is orthogonal to the null space, and they are orthogonal complements of one another,

$$C(A^T) = N(A)^\perp, \quad N(A) = C(A^T)^\perp.$$




Figure 1. Graphical representation of the Fundamental Theorem of Linear Algebra, Gil Strang, *Amer. Math. Monthly* **100**, 848-855, 1993.

Consideration of equality between sets arises in proving the above theorem. A standard technique to show set equality $A=B$, is by double inclusion, $A \subseteq B \wedge B \subseteq A \Rightarrow A=B$. This is shown for the statements giving the decomposition of the codomain \mathbb{R}^m . A similar approach can be used to decomposition of \mathbb{R}^n .

i. $C(A) \perp N(A^T)$ (column space is orthogonal to left null space).

Proof. Consider arbitrary $u \in C(A), v \in N(A^T)$. By definition of $C(A)$, $\exists x \in \mathbb{R}^n$ such that $u = Ax$, and by definition of $N(A^T)$, $A^T v = \mathbf{0}$. Compute $u^T v = (Ax)^T v = x^T A^T v = x^T (\mathbf{0}) = x^T \mathbf{0} = 0$, hence $u \perp v$ for arbitrary u, v , and $C(A) \perp N(A^T)$. \square

ii. $C(A) \cap N(A^T) = \{\mathbf{0}\}$ ($\mathbf{0}$ is the only vector both in $C(A)$ and $N(A^T)$).

Proof. (By contradiction, *reductio ad absurdum*). Assume there might be $b \in C(A)$ and $b \in N(A^T)$ and $b \neq \mathbf{0}$. Since $b \in C(A)$, $\exists x \in \mathbb{R}^n$ such that $b = Ax$. Since $b \in N(A^T)$, $A^T b = A^T (Ax) = \mathbf{0}$. Note that $x \neq 0$ since $x = 0 \Rightarrow b = 0$, contradicting assumptions. Multiply equality $A^T Ax = \mathbf{0}$ on left by x^T ,

$$x^T A^T Ax = \mathbf{0} \Rightarrow (Ax)^T (Ax) = b^T b = \|b\|^2 = 0,$$

thereby obtaining $b = 0$, using norm property 3. Contradiction. \square

iii. $C(A) \oplus N(A^T) = \mathbb{R}^m$

Proof. (iii) and (iv) have established that $C(A), N(A^T)$ are orthogonal complements

$$C(A) = N(A^T)^\perp, \quad N(A^T) = C(A)^\perp.$$

By Lemma 2 it results that $C(A) \oplus N(A^T) = \mathbb{R}^m$. \square

The remainder of the FTLA is established by considering $\mathbf{B} = \mathbf{A}^T$, e.g., since it has been established in (v) that $C(\mathbf{B}) \oplus N(\mathbf{A}^T) = \mathbb{R}^n$, replacing $\mathbf{B} = \mathbf{A}^T$ yields $C(\mathbf{A}^T) \oplus N(\mathbf{A}) = \mathbb{R}^m$, etc.